

Interpolation and approximation with polynomials

The theory (excerpts from lectures)

Polynomial interpolation

Values of some real function $y(x)$ at a finite set of *distinct* points are prescribed and we want to interpolate them by a polynomial $p(x)$, so that we can estimate intermediate values of the function $y(x)$. Let us denote by x_i , $i = 0, 1, 2, \dots, n$ the values of independent variable x and by y_i the prescribed values of the function $y(x)$ at x_i and summarize all given values in a table:

| | | | | | |
|----------|-------|-------|-------|---------|-------|
| x | x_0 | x_1 | x_2 | \dots | x_n |
| y | y_0 | y_1 | y_2 | \dots | y_n |

Generally, if we have $n + 1$ data points $[x_i, y_i]$, there is exactly one polynomial

$$p_n(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$$

of degree at most n going through all the data points. Its coefficients a_0, a_1, \dots, a_n are determined by $n + 1$ linear equations $p(x_i) = y_i$, $i = 0, 1, \dots, n$:

$$\begin{aligned} a_0 + a_1x_0 + a_2x_0^2 + \dots + a_nx_0^n &= y_0 \\ a_0 + a_1x_1 + a_2x_1^2 + \dots + a_nx_1^n &= y_1 \\ &\dots \\ a_0 + a_1x_n + a_2x_n^2 + \dots + a_nx_n^n &= y_n \end{aligned}$$

This can be expressed in matrix form as $\mathbf{Q}\mathbf{a} = \mathbf{y}$:

$$\mathbf{Q}\mathbf{a} \equiv \begin{bmatrix} 1 & x_0 & x_0^2 & \dots & x_0^n \\ 1 & x_1 & x_1^2 & \dots & x_1^n \\ \dots & \dots & \dots & \dots & \dots \\ 1 & x_n & x_n^2 & \dots & x_n^n \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \dots \\ a_n \end{bmatrix} = \begin{bmatrix} y_0 \\ y_1 \\ \dots \\ y_n \end{bmatrix} \equiv \mathbf{y} \quad (1)$$

The matrix is called Vandermonde matrix and for distinct values of x_i it is invertible, so the system (1) has unique solution for unknown vector of coefficients \mathbf{a} .

Note: The system (1) can be ill conditioned and solving it is not an efficient way how to compute the coefficients of the interpolation polynomial. Using *Lagrange interpolation polynomials* can be recommended instead.

Example 1

Consider the following table of data points

| | | | |
|----------|----|---|---|
| x | -1 | 1 | 2 |
| y | 8 | 4 | 5 |

Find the interpolation polynomial and use it for estimating the value at $x = 0.5$.

The solution

There are 3 data points, it follows that we are seeking the polynomial of the second order $p_2(x) = a_0 + a_1x + a_2x^2$ which has three unknown coefficients. Let us put together the equations (1):

$$\begin{aligned} a_0 + a_1 \cdot (-1) + a_2 \cdot (-1)^2 &= 8 \\ a_0 + a_1 \cdot 1 + a_2 \cdot 1^2 &= 4 \\ a_0 + a_1 \cdot 2 + a_2 \cdot 2^2 &= 5 \end{aligned}$$

or, in the matrix form

$$\begin{bmatrix} 1 & -1 & 1 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_n \end{bmatrix} = \begin{bmatrix} 8 \\ 4 \\ 5 \end{bmatrix} .$$

The solution is $a_0 = 5$, $a_1 = -2$ and $a_2 = 1$, so the interpolating polynomial is $p_2(x) = 5 - 2x + x^2$.

Value at 0.5 can be estimated as $p_2(0.5) = 5 - 2 \cdot 0.5 + 0.5^2 = 4.25$




Figure 1: Solution of Example 1: given data (represented by dots) interpolated by the polynomial of second degree.

Polynomial approximation

As before, we are given $n + 1$ discrete values of some real function $y(x)$. Now we want to *approximate* them by a polynomial $p(x)$, so that we can estimate intermediate values of the function. We do not require the polynomial going exactly *through* the data points, we are satisfied if it is just *close* to them. However, we usually require the order m of the polynomial to be rather small, often m is less than 3.

Note: The requirement of x_i to be distinct values is not needed here.

Method of least squares

In this approximation method, the term *close* to the data points means that we want to minimize the sum of squares of vertical distances $r_i = p(x_i) - y_i$ of the given points from the graph of the polynomial.

Let us start again with the requirement that the polynomial goes through all the given data points: $p(x_i) = y_i$, $i = 0, 1, \dots, n$. The result will be a system of equations similar to (1). Now the system matrix \mathbf{Q} is not square any more, however. Now it has m columns and n rows, m being much smaller than n , so that the system is over-determined and existence of solution generally cannot be expected:

$$\mathbf{Q} \mathbf{a} \equiv \begin{bmatrix} 1 & x_0 & x_0^2 & \dots & x_0^m \\ 1 & x_1 & x_1^2 & \dots & x_1^m \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ 1 & x_n & x_n^2 & \dots & x_n^m \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \dots \\ a_m \end{bmatrix} = \begin{bmatrix} y_0 \\ y_1 \\ \dots \\ \dots \\ y_n \end{bmatrix} \equiv \mathbf{y} \quad . \quad (2)$$

This problem can be fixed by solving a system of *normal equations* $\mathbf{Q}^T \mathbf{Q} \mathbf{a} = \mathbf{Q}^T \mathbf{y}$ instead:

$$\mathbf{Q}^T \mathbf{Q} \mathbf{a} \equiv \begin{bmatrix} n+1 & \sum_{i=0}^n x_i & \dots & \sum_{i=0}^n x_i^m \\ \sum_{i=0}^n x_i & \sum_{i=0}^n x_i^2 & \dots & \sum_{i=0}^n x_i^{m+1} \\ \sum_{i=0}^n x_i^2 & \sum_{i=0}^n x_i^3 & \dots & \sum_{i=0}^n x_i^{m+2} \\ \dots & \dots & \dots & \dots \\ \sum_{i=0}^n x_i^m & \sum_{i=0}^n x_i^{m+1} & \dots & \sum_{i=0}^n x_i^{2m} \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \dots \\ a_m \end{bmatrix} = \begin{bmatrix} \sum_{i=0}^n y_i \\ \sum_{i=0}^n x_i y_i \\ \sum_{i=0}^n x_i^2 y_i \\ \dots \\ \sum_{i=0}^n x_i^m y_i \end{bmatrix} \equiv \mathbf{Q}^T \mathbf{y} \quad (3)$$

which means that the error $\mathbf{r} = \mathbf{Q} \mathbf{a} - \mathbf{y}$ should be orthogonal to column space of the matrix \mathbf{Q} . The matrix $\mathbf{Q}^T \mathbf{Q}$ is symmetric positive semidefinite, and if the columns of \mathbf{Q} are linearly independent, then it is positive definite (and consequently invertible).

Note: $n + 1$ is the number of the given data points.

Example 1 - continued

Suppose there are 3 more data points added to the table:

| | | | | | | |
|----------|----|---|---|----|---|---|
| x | -1 | 1 | 2 | -1 | 0 | 2 |
| y | 8 | 4 | 5 | 7 | 4 | 6 |

Find the approximation polynomial of second degree.

The solution

There are more equations than unknown variables and the system (2) generally has no solution:

$$\mathbf{Q}\mathbf{a} = \begin{bmatrix} 1 & -1 & 1 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & -1 & 1 \\ 1 & 0 & 0 \\ 1 & 2 & 4 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_n \end{bmatrix} = \begin{bmatrix} 8 \\ 4 \\ 5 \\ 7 \\ 4 \\ 6 \end{bmatrix} = \mathbf{y} \quad .$$

System of normal equations (3):

$$\mathbf{Q}^T\mathbf{Q}\mathbf{a} = \begin{bmatrix} 6 & 3 & 11 \\ 3 & 11 & 15 \\ 11 & 15 & 35 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_n \end{bmatrix} = \begin{bmatrix} 34 \\ 11 \\ 63 \end{bmatrix} = \mathbf{Q}^T\mathbf{y} \quad .$$

The solution is $a_0 = 4.3158$, $a_1 = -1.8816$ and $a_2 = 1.25$, so the approximating polynomial is $p_2(x) = 4.3158 - 1.8816x + 1.25x^2$. The result is depicted in Figure 2 by magenta line. For comparison, the blue line is the polynomial interpolating the original three blue dots.




Figure 2: Solution of Example 1: original data (represented by blue dots) and added data (represented by red dots), all together approximated by the polynomial of second degree (magenta line). The original interpolating polynomial is blue.

Theorem 1: approximation by a polynomial using the least squares method

Suppose data points $x_i, y_i, i = 0, \dots, n$ are given so that the matrix $\mathbf{Q}^T \mathbf{Q}$ of the system (3) is invertible. Then the solution of (3) represents coefficients of the polynomial $p_m(x) = a_0 + a_1x + a_2x^2 + \dots + a_mx^m$ which minimizes the norm $\|\mathbf{r}\|_2$ of the vector of residuals $r_i = p_m(x_i) - y_i$ (called *quadratic deviation*) among all polynomials of degree at most m .

For better insight, two different ways of proving this theorem follow.

Proof 1 (using Calculus)

We want to minimize the quadratic deviation δ , or δ^2 :

$$\delta^2 \equiv \|\mathbf{r}\|_2^2 = \sum_{i=0}^n r_i^2 = \sum_{i=0}^n (p_m(x_i) - y_i)^2 \rightarrow \min .$$

The minimum of this sum, with respect to the unknown coefficients a_0, a_1, \dots, a_m of the polynomial $p_m(x)$, is found by setting the gradient of the function $\delta^2 \equiv S(a_0, a_1, \dots, a_m)$ to zero. Since it has $m+1$ variables, there are $m+1$ gradient equations

$$\begin{aligned} 0 &= \frac{\partial S}{\partial a_0} = \sum_{i=0}^n 2r_i \frac{\partial r_i}{\partial a_0} = \sum_{i=0}^n 2r_i \\ 0 &= \frac{\partial S}{\partial a_1} = \sum_{i=0}^n 2r_i \frac{\partial r_i}{\partial a_1} = \sum_{i=0}^n 2r_i x_i \\ 0 &= \frac{\partial S}{\partial a_2} = \sum_{i=0}^n 2r_i \frac{\partial r_i}{\partial a_2} = \sum_{i=0}^n 2r_i x_i^2 \\ &\dots \\ 0 &= \frac{\partial S}{\partial a_m} = \sum_{i=0}^n 2r_i \frac{\partial r_i}{\partial a_m} = \sum_{i=0}^n 2r_i x_i^m \end{aligned}$$

After substituting of $r_i = p_m(x_i) - y_i = a_0 + a_1x + \dots + a_mx^m - y_i$ and rearranging, system (3) is obtained.

Proof 2 is based on linear algebra and on following

Theorem 2:

Suppose \mathbf{A} is a symmetric and positive definite (*spd*) matrix, \mathbf{b} is a vector and $\mathbf{F}(\mathbf{x})$ is a quadratic functional $\mathbf{F}(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x} - 2\mathbf{x}^T \mathbf{b}$. Then $\mathbf{A}\bar{\mathbf{x}} = \mathbf{b} \Leftrightarrow \mathbf{F}(\bar{\mathbf{x}}) < \mathbf{F}(\mathbf{x}) \quad \forall \mathbf{x} \neq \bar{\mathbf{x}}$.

The proof of Theorem 1:

Let $\mathbf{A} = \mathbf{Q}^T \mathbf{Q}$, $\mathbf{b} = \mathbf{Q}^T \mathbf{y}$.

Let us express $\|\mathbf{r}\|_2^2$ as a quadratic functional similar to \mathbf{F} in Theorem 2:

$$\begin{aligned} \|\mathbf{r}\|_2^2 &= \|\mathbf{Q}\mathbf{a} - \mathbf{y}\|_2^2 = (\mathbf{Q}\mathbf{a} - \mathbf{y})^T (\mathbf{Q}\mathbf{a} - \mathbf{y}) = \mathbf{a}^T \mathbf{Q}^T \mathbf{Q} \mathbf{a} - \mathbf{y}^T \mathbf{Q} \mathbf{a} - \mathbf{a}^T \mathbf{Q}^T \mathbf{y} + \mathbf{y}^T \mathbf{y} \\ &= \mathbf{a}^T \mathbf{A} \mathbf{a} - \mathbf{b}^T \mathbf{a} - \mathbf{a}^T \mathbf{b} + \mathbf{y}^T \mathbf{y} = \mathbf{a}^T \mathbf{A} \mathbf{a} - 2\mathbf{a}^T \mathbf{b} + \|\mathbf{y}\|_2^2 = \mathbf{F}(\mathbf{a}) + \|\mathbf{y}\|_2^2 \end{aligned}$$

As $\|\mathbf{y}\|_2^2$ does not depend on \mathbf{a} , it can be seen that minimization of $\|\mathbf{r}\|_2^2 = \|\mathbf{Q}\mathbf{a} - \mathbf{y}\|_2^2$ with respect to \mathbf{a} is the same as minimization of $\mathbf{F}(\mathbf{a}) = \mathbf{a}^T \mathbf{A} \mathbf{a} - 2\mathbf{a}^T \mathbf{b}$ with respect to \mathbf{a} . From Theorem 2 it follows that this can happen only if \mathbf{a} is the solution of the system (3), i.e. $\mathbf{A}\mathbf{a} = \mathbf{b}$.

Note: If $\mathbf{A}\mathbf{a} = \mathbf{b}$, then \mathbf{r} and $\mathbf{Q}\mathbf{a}$ are perpendicular vectors:

$$\begin{aligned} \mathbf{r}^T \mathbf{Q} \mathbf{a} &= (\mathbf{Q} \mathbf{a} - \mathbf{y})^T \mathbf{Q} \mathbf{a} = \mathbf{a}^T \mathbf{Q}^T \mathbf{Q} \mathbf{a} - \mathbf{y}^T \mathbf{Q} \mathbf{a} = \mathbf{a}^T \mathbf{A} \mathbf{a} - \mathbf{b}^T \mathbf{a} = (\mathbf{A} \mathbf{a} - \mathbf{b})^T \mathbf{a} = 0 , \\ &\text{which means that } \mathbf{Q} \mathbf{a} \text{ is the orthogonal projection of } \mathbf{y} \text{ to the column space of } \mathbf{Q}. \end{aligned}$$

Illustration of Theorem 2 in R^2 (recapitulation from Calculus):

$$\begin{aligned} \mathbf{F}(\mathbf{x}) &= \mathbf{x}^T \mathbf{A} \mathbf{x} - 2 \mathbf{x}^T \mathbf{b} = \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} a & c \\ c & d \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + 2 \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = \\ &= a x_1^2 + 2c x_1 x_2 + d x_2^2 - 2x_1 b_1 - 2x_2 b_2 \end{aligned}$$

$$\text{grad } \mathbf{F}(\mathbf{x}) = \begin{bmatrix} \frac{\partial \mathbf{F}}{\partial x_1} \\ \frac{\partial \mathbf{F}}{\partial x_2} \end{bmatrix} = 2 \begin{bmatrix} a x_1 + c x_2 - b_1 \\ c x_1 + d x_2 - b_2 \end{bmatrix} = 2(\mathbf{A}\mathbf{x} - \mathbf{b})$$

it follows that \mathbf{F} can have one critical point only – the solution of $\mathbf{A}\mathbf{x} = \mathbf{b}$ (the solution is unique, because the matrix \mathbf{A} is spd and so it is invertible)

$$\mathbf{H} = \begin{bmatrix} \frac{\partial^2 \mathbf{F}}{\partial x_1^2} & \frac{\partial^2 \mathbf{F}}{\partial x_1 x_2} \\ \frac{\partial^2 \mathbf{F}}{\partial x_2 x_1} & \frac{\partial^2 \mathbf{F}}{\partial x_2^2} \end{bmatrix} = \mathbf{A}$$

\mathbf{A} is spd, so the critical point represents minimum and the graph of \mathbf{F} is an elliptic paraboloid oriented upwards.

Example 2

Consider the following table of seven data points

| | | | | | | | |
|----------|----|---|---|---|---|---|----|
| x | -1 | 0 | 0 | 1 | 1 | 2 | 4 |
| y | 5 | 6 | 5 | 7 | 6 | 8 | 11 |

Find the approximation polynomials of the first and the second order and the standard deviations.

The solution

The **first-order** polynomial is a line $p_1(x) = a_0 + a_1 x$, coefficients of which are determined by the linear system (3) with $m = 1$, $n = 6$:

$$\mathbf{A} = \begin{bmatrix} n+1 & \sum_{i=0}^n x_i \\ \sum_{i=0}^n x_i & \sum_{i=0}^n x_i^2 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} \sum_{i=0}^n y_i \\ \sum_{i=0}^n x_i y_i \end{bmatrix}$$

Let us extend the given table of data in order to compute the elements of the matrix and the right hand side

| | | | | | | | | sum |
|----------------------|----|---|---|---|---|----|----|------------|
| x | -1 | 0 | 0 | 1 | 1 | 2 | 4 | 7 |
| y | 5 | 6 | 5 | 7 | 6 | 8 | 11 | 48 |
| xy | -5 | 0 | 0 | 7 | 6 | 16 | 44 | 68 |
| x² | 1 | 0 | 0 | 1 | 1 | 4 | 16 | 23 |

which gives two linear equations for unknowns a_0 and a_1 :

$$\begin{bmatrix} 7 & 7 \\ 7 & 23 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \end{bmatrix} = \begin{bmatrix} 48 \\ 68 \end{bmatrix}.$$

The solution is $a_0 = 5.6071$, $a_1 = 1.25$ and the linear approximation of the given data is $p_1(x) = 5.6071 + 1.25x$.

$$\begin{aligned} \delta^2 &= \sum_{i=0}^7 (p_1(x_i) - y_i)^2 = \\ &= (4.3571 - 5)^2 + (5.6071 - 6)^2 + (5.6071 - 5)^2 + (6.8571 - 7)^2 + (6.8571 - 6)^2 + (8.1071 - 8)^2 + (10.6071 - 11)^2 = \\ &= (-0.6429)^2 + (-0.3929)^2 + 0.6071^2 + (-0.1429)^2 + 0.8571^2 + 0.1071^2 + (-0.3929)^2 = \\ &= 0.4133 + 0.1543 + 0.3685 + 0.0204 + 0.7346 + 0.0115 + 0.1544 = 1.8571 \\ \delta &= \sqrt{\delta^2} = 1.3628 \end{aligned}$$

The **second-order** polynomial is a parabola $p_2(x) = a_0 + a_1x + a_2x^2$, coefficients of which are determined by the linear system (3) with $m = 2$, $n = 6$:

$$\mathbf{A} = \begin{bmatrix} n+1 & \sum_{i=0}^n x_i & \sum_{i=0}^n x_i^2 \\ \sum_{i=0}^n x_i & \sum_{i=0}^n x_i^2 & \sum_{i=0}^n x_i^3 \\ \sum_{i=0}^n x_i^2 & \sum_{i=0}^n x_i^3 & \sum_{i=0}^n x_i^4 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} \sum_{i=0}^n y_i \\ \sum_{i=0}^n x_i y_i \\ \sum_{i=0}^n x_i^2 y_i \end{bmatrix}$$

Let us extend the table by three more rows:

| | | | | | | | | |
|-----------------------|----|---|---|---|---|----|-----|------------|
| | | | | | | | | sum |
| x | -1 | 0 | 0 | 1 | 1 | 2 | 4 | 7 |
| y | 5 | 6 | 5 | 7 | 6 | 8 | 11 | 48 |
| xy | -5 | 0 | 0 | 7 | 6 | 16 | 44 | 68 |
| x² | 1 | 0 | 0 | 1 | 1 | 4 | 16 | 23 |
| x²y | 5 | 0 | 0 | 7 | 6 | 32 | 176 | 226 |
| x³ | -1 | 0 | 0 | 1 | 1 | 8 | 64 | 73 |
| x⁴ | 1 | 0 | 0 | 1 | 1 | 16 | 256 | 275 |

which gives three linear equations for unknowns a_0 , a_1 and a_2 :

$$\begin{bmatrix} 7 & 7 & 23 \\ 7 & 23 & 73 \\ 23 & 73 & 275 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 48 \\ 68 \\ 226 \end{bmatrix}.$$

The solution is $a_0 = 5.5856$, $a_1 = 0.8313$, $a_2 = 0.1340$ and the quadratic approximation of the given data is $p_1(x) = 5.5856 + 0.8313x + 0.1340x^2$.

$$\delta^2 = (4.8883 - 5)^2 + (5.5856 - 6)^2 + (5.5856 - 5)^2 + (6.5509 - 7)^2 + (6.5509 - 6)^2 + (7.7842 - 8)^2 + (11.0548 - 11)^2 = 1.0819$$

$$\delta = \sqrt{\delta^2} = 1.0401$$

Another way, how to obtain the linear system above:

$$\mathbf{Q} = \begin{bmatrix} 1 & x_0 & x_0^2 \\ 1 & x_1 & x_1^2 \\ \dots & \dots & \dots \\ 1 & x_n & x_n^2 \end{bmatrix} = \begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 4 & 16 \end{bmatrix}, \quad \mathbf{Q}^T \mathbf{Q} = \begin{bmatrix} 7 & 7 & 23 \\ 7 & 23 & 73 \\ 23 & 73 & 275 \end{bmatrix}, \quad \mathbf{Q}^T \mathbf{y} = \begin{bmatrix} 48 \\ 68 \\ 226 \end{bmatrix}.$$