

(partial derivatives)

1. a) Compute all partial derivatives of $f(x, y) = \ln(2x - y) + 3x^3 - xy$ in a point $A = [1; 1]$.
b) Write a tangent line of the function in a cut $x \equiv 1$ in tangent point A.
2. a) Compute all partial derivatives of $f(x, y, z) = ze^{xyz}$ in a point $A = [0; 2; -1]$.
b) Write a tangent line of the function in a cut $y \equiv 2 \wedge z \equiv -1$ in tangent point A.
3. Verify that a function $u(x, y) = e^y(y^2 - x^2)$ is a solution of an equation

$$y \frac{\partial u}{\partial x} + x \frac{\partial u}{\partial y} = xu.$$

4. Verify that a function $u(x, t) = \sin(x - ta)$ (with parameter $a \in \mathbb{R}$) is a solution of an equation

$$\frac{\partial u}{\partial t} + a \frac{\partial u}{\partial x} = 0.$$

Differential and tangent (hyper-)plane

5. a) Write (total) differential of a function $f(x, y) = \frac{y}{x}$ in a point $A_0 = [2; 1]$.
b) Approximate the increment of the function between points A_0 and $A_1 = [2.1; 1.2]$ (i.e. $\Delta f = f(A_1) - f(A_0) = ?$)
6. By using the (total) differential, approximate the value of $f(0.97; 1.02; 0.99) = \frac{\sqrt[4]{0.97}}{1.02^3 \sqrt[3]{0.99}}$ (with 2 decimal places precision) hint: Use known value $f(1; 1; 1)$.
7. Given $f(x, y) = 3y^2 - 2x^2 + x$ and a point $T = [2; -1; ?]$.
a) Compute P.D. of the function in a point $[2; -1]$.
b) Find an equation of the tangent plane (τ) to the graph of the function at the point T .
8. Find an equation of the plane (τ) tangent to the graph of $f(x, y) = x \sin(x + y)$ at a point $T = [-1; 1; ?]$. Find also an equation of a line (ν) normal to the graph of f at point T .
9. a) Find an equation of the plane tangent to the graph of $f(x, y) = \ln(x + y)$ at a point $[1; 0; ?]$.
b) Use the result to approximate the functional value $f(A_1)$ in a point $A_1 = [1.1; 0.1]$.
10. Given $f(x, y) = 2x^2 - y^2$ and a plane $\sigma : 8x - 6y - z + 12 = 0$.
a) Find a plane (τ) tangent to the graph of f and parallel to the plane σ .
b) Find a line (ν) normal to the graph of f and normal to the plane σ .
11. Find an equation of the hyper-plane (τ) tangent to the graph of $f(x, y, z) = \ln(x^2 - y + 3z)$ at a point $T = [2; 1; 1; ?]$.
12. Given $f(x, y, z) = \ln(z + \sqrt{9 - x^2 - y^2})$,
a) Find Domain of definition of f and sketch it (at least in 2 cuts).
b) Find an equation of the hyper-plane (τ) tangent to the graph of f at a point $T = [0; 0; 1; ?]$.