

(directional derivative)

1. Given $f(x, y, z) = x^2 - 2y^2 - 3z^3 - 17$ and a point $A = [1; 1; 1]$, compute the directional derivative of f at point A in the direction given by a vector $\vec{s} = (1; 1; 1)$. What can you say about the function in this direction?
2. Given $f(x, y, z) = x^3y + \frac{x}{y^2} + 2z$ and a point $A = [-1; 1; 0]$,
 - a) determine the direction \vec{s} in which is the directional derivative at point A maximal.
 - b) Compute the derivative in this direction (\vec{s}) at point A .

Chain rule (derivatives of composite functions)

3. Given $f(u, v) = u^2 \ln v$, compute $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ when you know that $u(x, y) = \frac{x}{y}$ and $v(x, y) = 3x - 2y$.
4. Given unknown function $z(x, y) = f(u, v) = f(u(x, y), v(x, y))$ and functions $u(x, y) = x^2 - y^2$, $v(x, y) = e^{xy}$. Compute $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$ at point $A = [1; 2]$ when you know (from physics) that $\frac{\partial f}{\partial u}(A) = 1$ and $\frac{\partial f}{\partial v}(A) = 0$.

Implicitly defined functions

5. Given $F(x, y) = x^3 + y^3 - 6xy + 4$, verify that by the equation $F(x, y) = 0$ is implicitly defined function $y = f(x)$ near the point $A = [1; 1]$. Compute its derivative $\frac{df}{dx}$ at point $x_0 = 1$ and find an equation of tangent to the graph of $f(x)$.
6. Verify that by equation $x^3y + y^3x + x^2y - 3 = 0$ is implicitly defined function $y = f(x)$ near the point $A = [1; 1]$. Compute its derivative $\frac{df}{dx}$ at point $x_0 = 1$ and find an equation of normal to the graph of $f(x)$.
7. Given $F(x, y) = \sin(x + y) - y^2 \cos x$, verify that by the equation $F(x, y) = 0$ is implicitly defined function $y = f(x)$ in the neighborhood of the point $A = [\pi; 0]$. Compute its derivative $\frac{df}{dx}$ at point $x_0 = \pi$ and describe the behavior of $f(x)$ near point A (is it increasing or decreasing, how fast?).
8. Given $F(x, y) = x^3 + 2x^2y + y^4$ verify that by the equation $F(x, y) = 1$ is implicitly defined function $y = f(x)$ near the point $A = [2; -1]$. Compute the first and the second derivative of f at point $x_0 = 2$.